





# Inflammatory Bowel Disease Standardized Care Protocols

## 3. IBD PATIENTS ON BIOLOGIC – MAINTENANCE REGIME

### Nurse:

- 1. Have the patient contact the nurse to report HBI (<u>#1</u>) or Partial Mayo (<u>#2</u>) every 2 months. Record this value in the patient's chart. If the patient is flaring, refer to protocol 1. SUSPECTED IBD OUTPATIENT FLARE.
- 2. Review information collected during biologic administration, if available. If there are any adverse events or issues, review with the physician. If not, sign and have scanned to the patient chart.
  - a. Adalimumab Injection Visit Report and Patient Questionnaire (#4). The Patient Questionnaire is distributed to patients every 8 weeks via AbbVie Care.
  - b. BioAdvance<sup>©</sup> Infliximab Post-infusion Feedback Report (<u>#5</u>) and Patient Questionnaire (<u>#6</u>). The Patient Questionnaire is given to patients at the time of their infusion.
- 3. Send a message to support staff to make follow-up appointments for every 8 months, and to send out bloodwork requisitions (listed below).
- 4. Have the patient get his/her Infliximab (<u>#13</u> & <u>#14</u>) or Adalimumab (<u>#15</u> & <u>#16</u>) trough and antibody levels drawn immediately prior to week 30 and week 54 doses.

## Support Staff:

- 1. Arrange follow-up clinic visit every 8 months, after the initial 12-14-week follow-up.
- 2. Along with the appointment notification, send the patient the following lab requisitions:
  - Biologic maintenance labs for the patient to complete every 2 months (<u>#11</u>)
    - > Follow-up bloodwork (<u>#12</u>) to do before each appointment
- 3. Ensure that the patient has been scheduled for colonoscopy\* at 6-10 months, as per Protocol 2. IBD Patients on Biologic Induction Regime.

#### Physician:

- 1. Consider dose optimization at 6-10 month colonoscopy\* if not at mucosal healing (#13).
  - \* Flexible sigmoidoscopy for left-sided ulcerative colitis.