

Patient Information Sheet

Patient Instructions for Azathioprine (brand name = Imuran®)

1. Please read the patient information sheet on azathioprine before you begin taking your medicine.
2. Take azathioprine using the following schedule:
 - ⇒ 50 mg (1 tablet) daily for 2 weeks, then
 - ⇒ 100 mg (2 tablets) daily for 2 weeks, then
 - ⇒ 150 mg (3 tablets) daily for as long as your IBD health practitioner directs you
3. Your IBD health practitioner will request regular blood tests to assess white blood cell counts and liver function to determine if you are tolerating the medication. It is mandatory that you complete the required laboratory tests. Failure to do so could result in severe and permanent tissue injury. Please complete the blood tests according to the following schedule:
 - ⇒ Blood tests completed every week for 1 month
 - ⇒ After, complete blood tests every month for as long as you are taking azathioprine

Note: Your IBD health practitioner may alter the dose of your medication based upon the results of these tests.
4. It will take 12 to 16 weeks for azathioprine to reach its maximal effectiveness.
5. Some viral infections can cause the white blood cell count to decrease temporarily while you are taking azathioprine. If you have a viral infection like a cold, please visit your family doctor and request a white blood cell count to be completed. If the white blood cell count is low, the azathioprine will either be stopped or the dose lowered until the cell count returns to normal.

