

HOSPITALS

UNIVERSITY OF ALBERTA

Sander Veldhuyzen van Zanten, MD, PhD
DIRECTOR
AHS Edmonton Zone Section Head
Pernilla D'Souza, MD
Levinus A Dieleman, MD, PhD
Richard N Fedorak, MD
Brendan Halloran, MD
Vivian Huang, MD
Dina H J Kao, MD
Karen I Kroeker, MD
Adriana Lazarescu, MD
Farhad Peerani, MD
Gurpal S Sandha, MBBS
Richard Sultanian, MD
Karen Wong, MD
Sergio Zepeda-Gomez, MD

HEPATOLOGY

Vincent G Bain, MD
Juan Gonzalez- Abraldes, MD
Klaus S Gutfreund, MD
Constantine J Karvellas, MD
Mang M Ma, MD
SITE CHIEF
Andrew L Mason, MBBS
Aldo Montano-Loza, MD
Puneeta Tandon, MD
Winnie W S Wong, MD

Nurse Practitioners

Michelle Carbonneau, NP
Melissa Harriott, NP
Kimberly Newnham, NP

BASIC SCIENTIST/RESEARCH

Karen L Madsen, PhD

PUBLIC HEALTH SCIENCES

Karen J Goodman, PhD
Phil Jacobs, PhD

ROYAL ALEXANDRA

Lana Bistriz, MD
SITE CHIEF
James P Ferguson, MD
Leah M Gramlich, MD
Jennifer Jin, MD
Melissa Johnson, MD
Greg Lutzak, MD
Kata Matic, MD
Jill McDermid, MD
Amy Morse, MD
Daniel C Sadowski, MD
Dennis N Todoruk, MD
Clarence K W Wong, MD

HEPATOLOGY

Robert J Bailey, MD
Marilyn Zeman, MD

MISERICORDIA

Allen Lim, MD
Melanie Pinchbeck, MD
Lori M Stead, MD, PhD
Brennan M A Walters, MD
SITE CHIEF

GREY NUNS

Anand Bala, MD
Vijey Selvarajah, MD
Jesse Siffledeen, MD
SITE CHIEF
Connie M Switzer, MD

STURGEON

Naseem Hoque, MD
SITE CHIEF

Perianal Irritation Treatment

1. Sulcrate 4% in orabase

- 66.4 grams Sulcrate + 33.6 grams orabase = 100 grams 2:1 ratio

or

2. Sulcrate 4% Protective Cream

- Sucralfate 4 grams = Eucerin (Aquafor) 50 gram + Sterile Water 50 ml = 100 grams

or

3. Sulcrate 4% in Glaxal Base Cream

- Sucralfate 40 mg + Hydrophilic Base, Non-Greasy 1 gram

Usage

- Apply to the affected areas BID and PRN

Volume

- 30 gram tube